arXiv:1206.5702v1 [quant-ph] 25 Jun 2012

Fundamental Uncertainty as a Consequence of Locality

Oscar C. O. Dahlsten,[1, 2] Andrew J. P. Garner,[2] and Vlatko Vedral[1, 2]
[1]*Center for Quantum Technologies, National University of Singapore, Republic of Singapore*
[2]*Atomic and Laser Physics, Clarendon Laboratory,*
University of Oxford, Parks Road, Oxford OX13PU, United Kingdom

(Dated: March 8, 2013)

In this paper we introduce an operational physical locality principle, *branch locality*, similar to non-signaling but applicable to single systems. Branch locality restricts the dynamics that can be implemented in a spatial interferometer. We give a simple proof that any theory with state-changing dynamics either: (i) respects a particular uncertainty relation, or (ii) violates branch locality. Quantum theory and classical theory are in category (i), whereas category (ii) includes the theory known as box world. Box world is typically presented as reasonable because it satisfies the non-signaling condition, but here we demonstrate that if it is allowed non-trivial dynamics, it must violate a locality principle that any physical theory should respect. Our results hence present a compelling argument for the necessity of the uncertainty relation in quantum and post-quantum theories.

Introduction.—The Uncertainty principle, stipulating that certain observables cannot be specified simultaneously, is considered the cornerstone of quantum physics. It was emphasized by Heisenberg to be a departure point of quantum physics from the classical view which maintains that all properties of physical systems can be known at the same time [1]. Bohr used the quantum uncertainty to develop a broader view known as "complementarity" which, he emphasized, existed between different physical measurement scenarios [2]. The fact that quantum physics implies that a particle's position and momentum cannot be known simultaneously has been considered counterintuitive and there have been many unsuccessful attempts to go beyond it.

Here we show that uncertainty itself is required by a very simple principle we call "branch locality", which stipulates that any action in a spatial region having zero probability of containing a system cannot have any effect on the state of that system.

Our arguably simple proof makes use of the well-established convex framework for probabilistic theories [3–8], which has been used to show interesting results in the fields of information theory [6], statistical mechanics [9] and axioms for quantum theory [3, 7, 8]. As an operational framework, essentially any experiments yielding tables of data can be described in this way [3, 4]. A consequence of our results is that computation via spatial interferometry appears useless in any post-quantum theory without the uncertainty relation.

Branch locality is similar in spirit to a commonly used locality principle known as *non-signalling*: that acting on one system does not have any immediate observable impact on another system at a different (space-like) location. Quantum theory respects non-signalling as the state of a subsystem is invariant under local operations on the other subsystems. (Violating non-signalling is distinct from Bell-violation, which quantum theory, of course, does exhibit). It seems very unlikely that any post-quantum theory will violate non-signalling.

Branch locality (BL), in contrast to non-signalling, is applicable to *single* systems. We consider a system which may be found in certain spatial regions with some probability. We demand that if the system has no probability of being found in a particular region, then actions on that region cannot have an observable impact on the system. We call this innocent-sounding principle *branch* locality, because we are envisaging different branches of an interferometer, see Figure 1.



FIG. 1: *The Mach-Zehnder interferometer*: Branch locality states that if the (possibly post-quantum) photon is with probability 1 to be found in one of the branches of the interferometer, then operations on the other spatially disjoint branch cannot change the (operational) state of the photon.

In a standard quantum interferometer experiment one can change the state, and thus the statistics of some measurements, by local operations on the branches (e.g. by introducing an optical fibre detour in a branch). These operations respect BL. In contrast, we find that if a theory violates a particular instance of the quantum uncertainty relation, then *all* state transforms violate BL. In other words, a theory for such experiments must either be trivial or respect that uncertainty relation.

For alternative perspectives on interference in post-quantum theories see for example [10, 11].

Modelling an interferometer experiment beyond quantum theory.— In the case of quantum theory one may describe an ideal Mach-Zehnder interferometer (Figure 1) using a single qubit, i.e. a 2-dimensional Hilbert space. The state after the first beam-splitter can be expressed in the which-branch basis as $|\psi\rangle = c_{\rm up}|z_{\rm up}\rangle + c_{\rm low}|z_{\rm low}\rangle$. The observable giving the expected position corresponds to $z_{\rm up}|z_{\rm up}\rangle\langle z_{\rm up}| + z_{\rm low}|z_{\rm low}\rangle\langle z_{\rm low}|$ for some labels $z_{\rm up}$, $z_{\rm low}$ that we assign to the respective branches. It will here be convenient to label these ±1 respectively so that the observable is modelled by the Pauli-matrix $Z = |z_{\rm up}\rangle\langle z_{\rm up}| - |z_{\rm low}\rangle\langle z_{\rm low}|$. (The argument will also work for other labellings than ±1).

The state space of a qubit is the well-known Bloch sphere, see Figure 2. Here a state is represented by a real-numbered vector of expectation values: $[\langle X\rangle, \langle Y\rangle, \langle Z\rangle]^T$ where X and Y are the other two Pauli-matrices, and $\langle g\rangle = p(g = +1) - p(g = -1)$. Mixtures of states correspond to probabilistic (convex) combinations of these states, lying inside the sphere of pure states defined by



FIG. 2: State space: the quantum state space is the (Bloch) sphere. We also consider the possibility of other states outside the sphere. The case of the maximal cubic state space is an instance of the so-called 'box world'. We use branch locality to show that the only states allowed on the faces of the cube are those right in the centre of each face (where the sphere touches the cube).

$$\langle X\rangle^2 + \langle Y\rangle^2 + \langle Z\rangle^2 = 1. \quad (1)$$

The above equation constitutes an uncertainty relation; for example if $\langle Z\rangle = 1$ one must have $\langle X\rangle = \langle Y\rangle = 0$. The more familiar formulation in terms of standard deviations, that $\Delta X \Delta Z \geq \frac{1}{2}|\langle[X, Z]\rangle| = |\langle Y\rangle|$, is implied by Eq.1 (recall that $(\Delta g)^2 = \langle g^2\rangle - \langle g\rangle^2$).

The real vector used above amounts to an *operational* description of the state. One may now entertain the possibility of post-quantum states by associating them with points outside the sphere of pure quantum states. For example, the state $(1, 1, 1)^T$ is not allowed in quantum theory as it violates the uncertainty principle of Eq.1. We shall here a priori allow such states and later rule them out.

It will be crucial to our argument to consider transformations of states. These must take all allowed states to allowed states. They must also respect probabilistic mixtures: for a transform T acting on a mixture of two states, ν_1 and ν_2, we have $T\left(p_1\vec{\nu_1} + p_2\vec{\nu_2}\right) = p_1 T\left(\vec{\nu_1}\right) + p_2 T\left(\vec{\nu_2}\right)$. We show in the technical supplement, using established results on convex theories, that transforms are then real-numbered matrices, up to the subtlety that one should now add an extra component n to the state vector corresponding to the 'normalisation' of the state ($n = 1$ for normalised states).

We shall here consider states of the form

$$\vec{\nu} = [n, \langle Z\rangle, \langle X_1\rangle, \langle X_2\rangle, \langle X_3\rangle ...]^T ,$$

where we use the notation $\langle g\rangle = p(g = +1) - p(g = -1)$ (if a state is subnormalised, in which case this takes values in the range $-n$ to n). We shall consistently take $\langle Z\rangle$ to be associated with position, such that $Z = +1(-1)$ is associated with the upper (lower) branch (we show in the technical supplement that the argument does not depend on this labelling). The number of measurements used to specify the state is restricted to three in quantum theory, with X_1 and X_2 the Pauli matrices X and Y respectively. However we shall allow for a larger or smaller number than that.

A *theory* is specified by the set of states (which implicitly assumes a set of measurements and outcomes has been defined) and the allowed transforms. Apart from making general statements about theories, we shall refer to three concrete examples. The *quantum* qubit case has states represented as $\vec{\nu} = [n, \langle Z\rangle, \langle X\rangle, \langle Y\rangle]^T$. The allowed transforms consist of both reversible SO(3) transforms as well as linear transforms shrinking the sphere. Secondly, we shall call the case of a diagonal density matrix (s.t. $\langle X\rangle = \langle Y\rangle = 0$) the *classical* case, modelled as $\vec{\nu} = [n, \langle Z\rangle, 0, 0]^T$. Here any matrix preserving or shrinking the line of states is allowed. Finally, the maximal state space of all probabilistic mixtures of the corners $(n, \pm n, \pm n, ...)$ is known as the state-space of a single system in *box-world*. The special case of $(n, \pm n, \pm n)$ is termed a *gbit*[6]. This can (for n=1) be visualised as the square X-Z plane slice of the cube in Figure 2. Gbits are currently of great interest in the context of understanding whether there can be Popescu-Rohrlich (PR) boxes. These are hypothetical maximally Bell-violating systems, see [12] or the technical supplement. Roughly speaking, the gbits are to a PR-box what qubits are to a Bell state. More specifically, the pure states of a gbit, the corners of the maximal state space, are the conditional marginal states of a PR-box—see the technical supplement. Thus if a PR-box can exist then so can a gbit. In box-world the allowed transforms on single systems are taken to be any matrix that preserves or shrinks the state space (but our arguments will apply even if one is not so permissive with the tranforms).

Branch Locality.— Branch locality, as described in Fig 1, can now be formalised as an operational principle.

Definition 1 (Principle of *branch locality*)**.** *Physical actions on one region of space have no immediate effect on systems with no probability of being detected in that region. In particular, a system definitely in the upper branch such that its state is* $\vec{\nu}_{\rm up} =$

$[n, \langle Z\rangle, \langle X_1\rangle, \langle X_2\rangle ...]^T = [n, 1, \langle X_1\rangle, \langle X_2\rangle ...]^T$ *is left invariant by any transform on the lower branch:*

$$T_{\rm low}\vec{\nu}_{\rm up} = \vec{\nu}_{\rm up}. \tag{2}$$

(similarly $T_{\rm up}\vec{\nu}_{\rm low} = \vec{\nu}_{\rm low}$*.)*

We shall also impose a more obvious condition on operations at different branches. We take the state to be described by someone without access to outcomes of any measurements performed on the local branches. Local transformations T acting on the respective branches must then not alter the statistics associated with the Z measurement, i.e. if the transform takes $\langle Z\rangle$ to $\langle Z\rangle'$ then

$$\langle Z\rangle = \langle Z\rangle'. \tag{3}$$

Example 1: Box world (gbit).— Using the notation defined above, if we take a state to be in the upper branch with certainty it must have the form $\vec{\nu}_{\rm up} = [n, n, \langle X\rangle]^T$ (recall that n is the state normalisation with $n = 1$ for a normalised state).

Consider an operation on the lower branch. From the above considerations, Equation 2 and Equation 3 both hold. Recall moreover that the transform is a matrix. It follows from Equation 2 that

$$T_{\rm low}\vec{\nu}_{\rm up} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} n \\ n \\ \langle X\rangle \end{bmatrix} = \begin{bmatrix} (a+b)n + c\langle X\rangle \\ (d+e)n + f\langle X\rangle \\ (g+h)n + i\langle X\rangle \end{bmatrix} = \begin{bmatrix} n \\ n \\ \langle X\rangle \end{bmatrix}.$$

One can write down a similar expression for Equation 3. Consider the ranges of the different variables for gbits: n can take values in the range 0 to 1 and $\langle X\rangle$ as well as $\langle Z\rangle$ from -n to n. Note that even when $\langle Z\rangle = \pm 1$, $\langle X\rangle$ is free to take any value in the range $-n$ to n. Note also that $T_{\rm low}$ is independent of the state it acts on. It follows with a little work that

$$T_{\rm low} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Similarly by considering $\langle Z\rangle = -1$ we have $T_{\rm up} = \mathbb{1}$.

Thus no non-trivial operation satisfying branch locality can be applied on either branch. However, there are clearly transforms leaving Z invariant that are not of this trivial form, e.g. $T^* = {\rm diag}(1, 1, -1)$. This transform if allowed would provide non-trivial interference. However, we have shown that T^* cannot be associated with a local operation on a branch! If we assume all box-world states can be realised in the interferometer, and any non-trivial transform can be performed, we must conclude that *box world violates branch locality.*

Uncertainty principle.—The above proof for a gbit that all non-trivial dynamics violate branch locality does not carry through in quantum theory. The proof makes use of that fact that, for gbits, when $\langle Z\rangle = \pm 1$, $\langle X\rangle$ is still free to take all possible values. This is a violation of the quantum uncertainty principle and a key difference between box world and quantum theory. We now argue that *any* convex framework theory violating such an uncertainty relation (defined below) has no state-changing dynamics respecting branch locality.

We consider a set of maximally independent measurements Z and $X_1 \ldots X_n$, all taking values -1 and 1. (We call a set of measurements $\{g_i\}_i$ *maximally independent* if we can write $\vec{\nu} = \sum_i \langle g_i\rangle \vec{v}_i$ and $\vec{v}_i \cdot \vec{v}_j = \delta_{ij}$.)

Definition 2 (Uncertainty principle for maximally independent measurements)**.** *If* $|\langle Z\rangle| = 1$ *then* $\langle X_i\rangle = 0\,\forall i$.

The uncertainty principle is said to be totally violated *if the set of states where* $|\langle Z\rangle| = 1$ *includes at least one state where* $\langle X_1\rangle \neq 0$*, one state where* $\langle X_2\rangle \neq 0$*, etc.*

Theorem 1. *In theories that satisfy branch locality, the transforms T on local branches (thus respecting Eq. 3) must have the form:*

$$T = \mathbb{1}$$

except when the uncertainty principle is upheld.

The proof, which follows similar lines to the example above in a more complicated setting, is in the technical supplement.

Example A: Qubit.— For pure states of a qubit $\langle X\rangle^2 + \langle Y\rangle^2 + \langle Z\rangle^2 = 1$. Thus when $\langle Z\rangle = \pm 1$ both $\langle X\rangle = 0$ and $\langle Y\rangle = 0$. Thus, labelling the measurements $X_1 = X$ and $X_2 = Y$, we see that a qubit satisfies the uncertainty principle of Definition 2. Theorem 1, does accordingly *not* imply that the transforms in this case have to be $\mathbb{1}$. Instead, one can see that imposing BL, in particular Equations 3 and 2, implies that

$$T = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & a & b \\ 0 & 0 & c & d \end{bmatrix}$$

for both the upper and lower branches. This confirms what is well-known, namely that quantum theory can describe dynamics other than $\mathbb{1}$ in an interferometer.

Example B: Box world.—Box world, even beyond the 2-dimensional gbit case, violates the uncertainty principle. Thus from Theorem 1, the only operation that may be applied without violating branch locality is the trivial element $T = \mathbb{1}$. In other words, all box-world transforms violate BL. Note that this statement does not only apply to box-world, but to any state space with more states on the faces of the outer cube than those right in the centres of the faces.

Example C: Classical.—The classical case (also defined above) is a subspace within quantum theory, and thus surprisingly respects the uncertainty relation. Accordingly Theorem 1 does not have a bearing on transformations in this case. Nevertheless one sees the transformations are trivial as there is only one degree of freedom, Z, and from Equation 3, it is conserved.

Example D: Which branch measurement.—We now give an interesting example of a transform which should respect branch locality in any theory: a measurement of which branch the particle is in. It is not obvious a priori that this should respect branch locality as it appears to be a multi-branch operation. Suppose however that one places a (non-destructive) detector in the lower path only. Physically this is an operation on the lower branch only, and the associated transform of the state should accordingly leave states associated with the upper branch invariant, i.e. it should respect branch locality. Moreover Eq. 3 must be respected. Accordingly, our results can be applied here to imply that the state transformation associated with the detector has to be $\mathbb{1}$, *unless* the uncertainty relation is respected. This suggests a close link between our results and measurement disturbance (as a disturbing transform cannot equal $\mathbb{1}$).

Consequences for computation.— As computers are physical objects, the laws of nature dictate the limits of computation. In particular, a computation has to be performed as a dynamical evolution of a physical system. Our results can be interpreted as saying that computation using a Mach-Zehnder interferometer is trivial unless the uncertainty relation holds. Whilst this is a very particular experimental setting, it stands out as being the original setting in which quantum computation was conceived. The Deutsch-Jozsa algorithm [13] arises naturally by considering what one can do with a quantum system in an interferometer. In [14] it was argued more generally that key quantum algorithms can be viewed as a three stage interferometer experiment: (i) prepare a superposition of different branches, (ii) apply different phase-shifts to different branches, and (iii) bring the branches together and make a measurement, yielding information about the phase shifts that were done. The apparent ability to prepare and individually address several inputs in stages (i) and (ii) is called quantum parallelism [13] (distinct from classical parallel computation).

Our result accordingly hints that an uncertainty relation is required to achieve this parallelism, particularly if the results for spatial degrees of freedom can be applied to all degrees of freedom. It suggests that parallelism may not even be a sensible concept in a spatially disjoint system without uncertainty. This directs the search for post-quantum theories with stronger computational power to those that respect the uncertainty relation, such as ‘systems with limited information content’ [15].

It may seem counterintuitive that the *restriction* of uncertainty leads to better computation, but this is because branch locality places a joint restriction on the states and transforms, such that the restriction on transforms is weakened by restricting the set of states.

It is already known that box world has very poor dynamics [6, 16], but the branch locality principle has more dramatic implications than previous results. It rules out any non-trivial normalisation-preserving dynamics, whether reversible or not, and by non-trivial we mean any transform that is not the identity (whereas the word ‘trivial’ in the title of [16] refers to the lack of correlating interactions). Moreover we showed the same restriction holds for *any* convex theory on bits not satisfying the above uncertainty relation.

Relation to other principles.—The principle is, as already mentioned, very similar to non-signalling; the two principles seem to be instances of some wider notion of locality. More subtly they can also both be viewed as *defining* what it means to have an operation on a subsystem or, in the case of branch locality, an operation on a branch. It would seem impossible to talk about notions like systems localised in space without these principles. We think they are both equally fundamental and unlikely to be violated in a post-quantum theory. However, standard non-signalling does not rule out box world, which has been designed to respect it. It is surprising that another locality principle does [6].

There are also many interesting information theoretic principles one may compare and relate branch-locality to. The one-bit principle, as borne out e.g. in [7, 17, 18], has a very similar mathematical implication, but with a radically different motivation. The principle, in one of its many similar formulations, states that an elementary system (a system with 2 perfectly distinguishable states) can only give a definite result in one specific measurement. The mathematical formulation of this in [18] is actually very close to what we here call the uncertainty principle for maximally independent measurements. Our argument accordingly supports the one-bit principle.

Outlook.— Whilst the uncertainty principle we consider above is a very significant restriction, the full quantum uncertainty relation also restricts states where none of the measurements are deterministic. It would be interesting to investigate possible extensions of branch locality to include such cases. For example one would expect that states with little weight on one branch can only be slightly changed by operations on that branch. Alternatively one may investigate combining branch locality with some other fundamental principle(s), as part of a program of exploring which convex theories are consistent with space-time.

As there is a relation between the uncertainty relation and the amount of Bell violation in a theory [19, 20], one may hope to link Tsirelson’s upper bound on quantum correlations with branch-locality.

We have refrained from claiming that branch locality applies to other degrees of freedom than position, but we anticipate that it is (this is actually why we call it branch locality rather than path locality which could otherwise have been more natural).

Acknowledgements.—We gratefully acknowledge discussions and correspondence (in chronological order) with Yoshifumi Nakata, Mio Murao, Markus Müller, Časlav Brukner, Mehdi Ahmadi, Anton Zeilinger and Matt Pusey as well as funding from the National Research Foundation (Singapore), the Ministry of Education (Singapore), the EPSRC (UK), the Templeton Foundation and the Leverhulme Trust. OD was regularly visiting Imperial College whilst undertaking this research.

---

[1] W. K. Heisenberg, *The physical principles of the quantum theory* (Dover, New York, NY, 1930).
[2] N. Bohr, American Journal of Physics **26** (1958).
[3] L. Hardy, (2001), arXiv:quant-ph/0101012.
[4] P. Mana, arXiv:quant-ph/0305117v3 (2003).
[5] H. Barnum, J. Barrett, M. Leifer, and A. Wilce, Phys. Rev. Lett. **99**, 240501 (2007).
[6] J. Barrett, Phys. Rev. A **75**, 032304 (2007).
[7] B. Dakic and C. Brukner, Deep Beauty(...) Ed. H. Halvorson, CUP , 365 (2011).
[8] L. Masanes and M. P. Müller, NJP **13**, 063001 (2011).
[9] M. P. Müller, O. C. O. Dahlsten, and V. Vedral, arXiv:1107.6029v1 [quant-ph] (To appear in Comm. Math. Phys.) (2011).
[10] R. Sorkin, Mod. Phys. Lett. A **9**, 3119 (1994).
[11] C. Ududec, H. Barnum, and J. Emerson, Found Phys **41**, 15 (2011), arXiv:1003.5005.
[12] S. Popescu and D. Rohrlich, Found. Phys. **24**, 379 (1994).
[13] D. Deutsch and R. Jozsa, Proc. R. Soc. Lond. A. **439**, 553 (1992).
[14] R. Cleve, A. Ekert, C. Macchiavello, and M. Mosca, Proc. R. Soc. Lond. A. **454**, 339 (1998).
[15] T. Paterek, B. Dakic, and C. Brukner, NJP **12**, 053037 (2010).
[16] D. Gross, M. Müller, R. Colbeck, and O. C. O. Dahlsten, Phys. Rev. Lett. **104**, 080402 (2010).
[17] A. Zeilinger, Found.Phys. **29** (1999).
[18] C. Brukner and A. Zeilinger, Found. Phys. **39**, 677 (2009).
[19] G. verSteeg and S. Wehner, QIC **9**, 0801 (2009).
[20] J. Oppenheim and S. Wehner, Science 19 **330**, 1072 (2010), arXiv:1004.2507.

Technical supplement

Introduction to the general probabilistic theory (GPT) framework

We firstly introduce the key concepts of the operationalist approach we will use: *the framework for general probabilistic theories (GPT)* also known as the convex framework. For a more detailed description of the framework see e.g. [3, 8]. The framework is operational in the sense that essentially any experiment producing a data table can be described in this way [3, 4]. For readers familiar with quantum theory it can also be helpful to think of the GPT framework as a generalisation of quantum theory. In quantum theory there is a system which is prepared in a state ρ, determined by the preparation in question. There is a set of measurements one may do, each represented by a set of projection operators $\{\Pi_i\}_{i=1}^{\dim\mathcal{H}}$ (or more generally POVM elements). The operationally significant quantities, the probabilities of given outcomes are given by $p_i = Tr(\rho\Pi_i)$. Viewed more abstractly, the state is a vector $\vec{\rho}$ in the vector space of Hermitian operators. A projection operator is also such a vector, $\vec{\Pi}$ say. In other words we may pick a basis of the vector space and write $\vec{\rho} = \sum_i \xi_i \vec{e_i}$ and $\vec{\Pi} = \sum_j \nu_j \vec{e_j}$. (Examples of such bases are the Pauli operators and the pure state basis given by Hardy in [3]). Note that the coefficients of these expansions are *real*, so this is termed a real vector space. $Tr(\rho\Pi)$ is then the Hilbert-Schmidt inner product (for Hermitian matrices) which we may write as $\langle\vec{\rho},\vec{\Pi}\rangle$.

If the basis elements are chosen so that they are orthogonal with respect to the norm, and all having the same inner product c with themselves, we see that $\langle\vec{\rho},\vec{\Pi}\rangle = \vec{\xi}\cdot\vec{\nu}c$ where the right-hand-side is the standard Euclidean norm. Thus we may represent a quantum state, measurements on it, and the resulting probabilities in terms of real vectors and the Euclidean norm.

In the framework we more generally represent the state of a system $\vec{s}$ as a real vector, and the measurement-outcome pairs, called 'effects' for historical reasons, as real vectors $\vec{e}$ (such a vector could for example in the quantum case be associated with X=+1, where X is the Pauli X). The probability of the outcome associated with a given $\vec{e}$ is given by $\vec{s}\cdot\vec{e}$. Part of specifying a given theory is specifying which states and effects are allowed. All convex combinations (mixtures) of allowed states are always allowed, written as $\vec{s} = \sum_i p_i \vec{s_i}$ (hence the name 'convex framework'). A state is said to be *pure* if it is not a non-trivial mixture of other states, otherwise it is called *mixed*.

Transformations are represented as real matrices acting on the state vector (following from the requirement of respecting mixtures, see e.g. [3]). They must take all allowed states to allowed states, but there may be further restrictions specified for a given theory in the framework. A transformation T is termed *reversible* if its inverse T^{-1} is also an allowed operation.

As well as quantum theory and theories contained therein (such as classical probability theory), one may also formulate a theory called *box world* in this way. Box world contains all states that do not violate non-signalling (that the reduced state of one system is invariant under operations on another) [6]. The standard version of box world assumes there are only two binary outcome measurements under considerations. We label these X and Z and the outcomes ±1 in analogy with quantum theory. A normalised state can then, as discussed below, be represented as $\vec{\nu} = [\langle X\rangle\ \langle Z\rangle]^T$ and is any mixture of the four extremal states $[\pm 1\ \ \pm 1]^T$. The most general box worlds are m-in n-out box worlds, which mean that one selects one measurement setting from m possible settings and obtain n-valued outcomes. In particular, the 3-in 2-out box world is the most analogous to a qubit in quantum theory.

We shall assume throughout that if a given $\vec{\nu}$ is allowed, the same state with one or more of the entries replaced by 0 is also allowed (an entry being 0 corresponds to that observable being completely unknown).

Expectation value representation

We shall here find it makes the notation very simple to focus on the expectation values of measurements, along the lines of the above example for box-world. In what follows we define a representation of states in terms of expectation values and relate it to the more standard representation in terms of probabilities, including showing that the transforms are matrices also in the new representation.

Consider the case of states described in terms of two fiducial measurements with binary outcomes (measurements are called fiducial if their statistics are sufficient to determine the state). We shall here only consider measurements with binary outcomes (because we shall be interested in interferometers with two branches), but we will later generalise to many such measurements. Such outcomes need not be normalised, but we require the sum of probabilities of both measurements to be equal.

Consider the *probability* representation of a state [6]:

$$\vec{s} := \left[\begin{array}{c} p(X=+1) \\ p(X=-1) \\ \hline p(Z=+1) \\ p(Z=-1) \end{array}\right].$$

The following is the alternative (normalisation-including) *expectation* value representation:

$$\vec{\nu} := \begin{bmatrix} n \\ p(X=+1)-p(X=-1) \\ p(Z=+1)-p(Z=-1) \end{bmatrix} = \begin{bmatrix} n \\ \langle X\rangle \\ \langle Z\rangle \end{bmatrix},$$

where the normalisation $n = 1$ if the state is normalised and $n < 1$ if it is subnormalised. (For a subnormalised state we still use the notation $\langle g\rangle := p(g=+1)-p(g=-1)$, implying that $-n\leq\langle g\rangle\leq n$)

If a transform T acts as a matrix on the state-vector in the probability representation, as it should if it respects mixtures, is it also a matrix in the expectation value picture? Suppose for the sake of argument that (i) There exists a fixed matrix M such that $\vec{\nu}=M\vec{s}$ for all states, (ii) The effective inverse matrix M^{-1} also exists satisfying $M^{-1}M\vec{s}=\vec{s}\;\forall\vec{s}$. Then we can write

$$MT\vec{s}=MTM^{-1}M\vec{s}=MTM^{-1}\vec{\nu}=\vec{\nu'},$$

where $\vec{\nu'}$ is the expectation representation state after the transform. Thus we see that *if the two assumptions above hold then the state transforms by a matrix also in the expectation value picture.* Moreover these two assumptions do hold here, with

$$M=\begin{bmatrix}1/2&1/2&1/2&1/2\\1&-1&0&0\\0&0&1&-1\end{bmatrix},\;M^{-1}=\begin{bmatrix}1/2&1/2&0\\1/2&-1/2&0\\1/2&0&1/2\\1/2&0&-1/2\end{bmatrix}.$$

The above argument naturally generalises to more measurements.

Not also that one could have used a different label for the positions in the probability picture, i.e. not ±1 and then mapped that into the expectation value picture using the same matrix as above. In this sense our argument does not depend on how one labels the two positions.

A. PR-boxes and gbits

We now briefly describe the connection between PR-boxes and gbits. This subsection is included for completeness and does not contain new results, see in particular [6] for essentially the same statements.

In the probability representation a general gbit state is a mixture of four pure states:

$$\vec{s}_{\rm gbit}:=p_1\left[\begin{array}{c}1\\0\\\hline1\\0\end{array}\right].+p_2\left[\begin{array}{c}1\\0\\\hline0\\1\end{array}\right].+p_3\left[\begin{array}{c}0\\1\\\hline1\\0\end{array}\right].+p_4\left[\begin{array}{c}0\\1\\\hline0\\1\end{array}\right].$$

Thus under the assumptions that (i) the four pure states are allowed and (ii) any probabilistic mixtures of states are allowed, then all such states are allowed.

Now consider a PR-box. This involves two systems, labelled A and B here. The state can be represented as

$$\vec{s}:=\left[\begin{array}{c}p(X_A=+1,X_B=+1)\\p(X_A=+1,X_B=-1)\\p(X_A=-1,X_B=+1)\\p(X_A=-1,X_B=-1)\\\hline p(X_A=1,Z_B=+1)\\p(X_A=+1,Z_B=-1)\\p(X_A=-1,Z_B=+1)\\p(X_A=-1,Z_B=-1)\\\hline p(Z_A=+1,X_B=+1)\\p(Z_A=+1,X_B=-1)\\p(Z_A=-1,X_B=+1)\\p(Z_A=-1,X_B=-1)\\\hline p(Z_A=+1,Z_B=+1)\\p(Z_A=+1,Z_B=-1)\\p(Z_A=-1,Z_B=+1)\\p(Z_A=-1,Z_B=-1)\end{array}\right]=\left[\begin{array}{c}\frac{1}{2}\\0\\0\\\frac{1}{2}\\\hline\frac{1}{2}\\0\\0\\\frac{1}{2}\\\hline\frac{1}{2}\\0\\0\\\frac{1}{2}\\\hline0\\\frac{1}{2}\\\frac{1}{2}\\0\end{array}\right]$$

Now consider for example the case where a measurement of X_A is performed, and one finds $X_A=+1$. Then we see the state of B, called the *conditional marginal state*, must be

$$\vec{s_B}=\left[\begin{array}{c}p(X_B=+1)\\p(X_B=-1)\\\hline p(Z_B=+1)\\p(Z_B=-1)\end{array}\right]=\left[\begin{array}{c}1\\0\\\hline1\\0\end{array}\right],$$

i.e. the conditional marginal state is one of the four pure states of the gbit state space. One can see by considering $X_A=-1$, $Z_A=\pm1$ that the other three pure states can be similarly prepared. Thus under the assumption that a PR-box is allowed, and that mixtures of all states are allowed, then the most general gbit-state is allowed.

B. Proof of main theorem

Theorem. *In theories that satisfy branch locality, the transforms T on local branches must have the form:*

$$T=\mathbb{1}$$

except when the uncertainty principle is upheld.

Proof. For T to leave the statistics of Z invariant, it must satisfy, for all $\langle Z\rangle$ and n

$$T\vec{\nu}=\begin{bmatrix}r&n_0&n_1&\dots&n_n\\z_{\rm r}&z_0&z_1&\dots&z_n\\x_{0{\rm r}}&x_{00}&x_{01}&\dots&x_{0n}\\\vdots&\vdots&\vdots&&\vdots\\x_{n{\rm r}}&x_{n0}&x_{n1}&\dots&x_{nn}\end{bmatrix}\begin{bmatrix}n\\\langle Z\rangle\\\langle X_1\rangle\\\vdots\\\langle X_n\rangle\end{bmatrix}=\begin{bmatrix}n\\\langle Z\rangle\\\langle X_1\rangle'\\\vdots\\\langle X_n\rangle'\end{bmatrix}.$$

This gives a set of simultaneous equations:

$$\begin{aligned}rn+n_0\langle Z\rangle+n_1\langle X_1\rangle+\ldots+n_n\langle X_n\rangle&=n, &(4)\\z_{\rm r}n+z_0\langle Z\rangle+z_1\langle X_1\rangle+\ldots+z_n\langle X_n\rangle&=\langle Z\rangle, &(5)\\x_{i{\rm r}}n+x_{i0}\langle Z\rangle+x_{i1}\langle X_1\rangle+\ldots+x_{in}\langle X_n\rangle&=\langle X_i\rangle'.&(6)\end{aligned}$$

These must hold for all allowed n, $\langle Z\rangle$ and $\langle X_1\rangle \dots \langle X_n\rangle$. As 0 is always a possible expectation value (we can always choose to forget the outcome of a measurement), we can set $\langle Z\rangle = 0$ and $\langle X_i\rangle = 0\,\forall i$, and so Equation 4 shows that $r = 1$. We can choose non-zero $\langle Z\rangle$ and see from Equation 5 that $z_0 = 1$, and from Equation 4 that $n_0 = 0$. We can also choose to set all but one $\langle Z\rangle$, $\langle X_i\rangle = 0$, so that Equation 5 implies that each $z_i = 0$ for $i = 1 \dots n$, and Equation 4 that each $n_i = 0$ for $i = 0 \dots n$, giving us the form of the top two rows of the transform.

Moreover, for T to respect branch locality, taking Z to be the position measurement, we can set $\langle Z\rangle = n$ and $\langle X_i\rangle' = \langle X_i\rangle$ to give the set of equations:

$$x_{i\mathrm{r}}n + x_{i0}n + x_{i1}\langle X_1\rangle + \ldots + x_{in}\langle X_n\rangle = \langle X_i\rangle. \tag{7}$$

Again, we can choose to set all $\langle X_i\rangle = 0$, and so Equation 7 will reduce to $x_{i\mathrm{r}} + x_{i0} = 0\,\forall i$.

When uncertainty is totally violated we are free to set all $\langle X_i\rangle = 0$ except one $\langle X_j\rangle \neq 0$, and from Equation 7 we see that $x_{ji} = \delta_{ji}$ where δ_{ji} is the Kronecker delta. Furthermore, normalisation constraints will require $x_{i\mathrm{r}}, x_{i0} = 0$. Thus, for total violation of the uncertainty principle, the only operation allowed is $T = \mathbb{1}$.

The case where the uncertainty principle is partially violated (i.e. there exists at least one non-zero value of $\langle X_i\rangle$ for some i), then it can also be shown that that the sub-matrix linking measurements without an uncertainty constraint is the identity. The proof is nearly identical to the case for total violation and so is omitted.

□